arXiv:hep-th/0404128v1 19 Apr 2004

Topological Charge of ADHM Instanton on $\mathbf{R}^2_{\rm NC} \times \mathbf{R}^2$

Yu Tian
Institute of Theoretical Physics, Chinese Academy of Sciences
P. O. Box 2735, Beijing 100080, China
`ytian@itp.ac.cn`

February 1, 2008

Abstract

We have calculated the topological charge of $U(N)$ instantons on non-degenerate noncommutative space time to be exactly the instanton number k in a previous paper [1]. This paper, which deals with the degenerate $\mathbf{R}^2_{\rm NC} \times \mathbf{R}^2$ case, is the continuation of [1]. We find that the same conclusion holds in this case, thus complete the answer to the problem of topological charge of noncommutative $U(N)$ instantons.

In a previous article [1], we have explicitly shown that the topological charge of $U(N)$ instantons on non-degenerate noncommutative space time[1] $\mathbf{R}^4_{\mathrm{NC}}$ is equal to the instanton number k, which appears in the noncommutative ADHM construction [6] as the dimension of the vector space V. There we left the problem of degenerate noncommutative parameter θ. We mentioned that our method ceased to work in the limit of $\theta_2 \to 0$ because of the loss of a Fock space representation for z_2 and $\bar{z}_2$.[2]

In this degenerate limit, the noncommutative space time becomes a direct product $\mathbf{R}^2_{\mathrm{NC}} \times \mathbf{R}^2$. It is well known that field theories on $\mathbf{R}^4_{\mathrm{NC}}$ suffer from a common problem of non-unitarity. But the problem does not happen on $\mathbf{R}^2_{\mathrm{NC}} \times \mathbf{R}^2$ if we require a time in the commutative part. For this reason, $\mathbf{R}^2_{\mathrm{NC}} \times \mathbf{R}^2$ may be of more physical relevance than $\mathbf{R}^4_{\mathrm{NC}}$. Literature has also shown that nonsingular ADHM instanton solutions can be constructed on $\mathbf{R}^2_{\mathrm{NC}} \times \mathbf{R}^2$ without involving any projected states [7, 8].

In the following paragraphs, we will analytically calculate the topological charge of $U(N)$ instantons on $\mathbf{R}^2_{\mathrm{NC}} \times \mathbf{R}^2$. As the continuation of [1], this paper will be brief about the relevant background and rely on some results in [1].

First let us recall the commutator of complex coordinates:

$$[\bar{z}_1, z_1] = \theta_1 = \zeta, \quad [\bar{z}_2, z_2] = \theta_2 = 0. \tag{1}$$

Now we introduce a Fock space representation for z_1 and $\bar{z}_1$:

$$\begin{aligned} z_1|n\rangle &= \sqrt{\zeta}\sqrt{n+1}|n+1\rangle, && (2)\\ \bar{z}_1|n\rangle &= \sqrt{\zeta}\sqrt{n}|n-1\rangle. && (3) \end{aligned}$$

Then the integration on $\mathbf{R}^2_{\mathrm{NC}} \times \mathbf{R}^2$ can be expressed as

$$\int d^4x\, \mathcal{O}(x) = \zeta\pi \sum_{n=0}^{\infty} \int d^2x\, \langle n|\mathcal{O}(x)|n\rangle, \tag{4}$$

where $d^2x = dx^3 dx^4$ [8].

Using the Corrigan's identity [9], we have the following expression of topological charge of (ASD) ADHM instanton on $\mathbf{R}^2_{\mathrm{NC}} \times \mathbf{R}^2$ with instanton number k [1]:

$$Q = \frac{1}{16\pi^2} \int d^4x\, \mathrm{Tr}_N (F_{mn} F_{mn}) \tag{5}$$

---

[1] For noncommutative geometry and field theory, please see [2, 3, 4, 5].

[2] Here we follow the notation of [1].

$$\begin{aligned} &= \frac{1}{16\pi^2}\int d^4x\,(8\partial\bar{\partial}+2\partial_\mu\partial_\mu)\mathrm{Tr}[b^\dagger(2-\Delta f\Delta^\dagger)bf] && (6)\\ &= \frac{\zeta}{\pi}\sum_{n=0}^{\infty}\int d^2x\,\langle n|(\frac{1}{2}\partial\bar{\partial}+\frac{1}{8}\partial_\mu\partial_\mu)\mathrm{Tr}[(2-\Delta' f\Delta'^\dagger)f]|n\rangle, && (7)\end{aligned}$$

where $\partial \equiv \partial_{z_1}$, $\bar{\partial} \equiv \partial_{\bar{z}_1}$, $\mu = 3, 4$, and $\Delta' \equiv b^\dagger\Delta$ [10].

Similarly as in [1], we have the truncated summation

$$\sum_{n=0}^{N}\langle n|\partial\bar{\partial}\mathcal{O}(x)|n\rangle = \frac{N+1}{\zeta}(\langle N+1|\mathcal{O}|N+1\rangle - \langle N|\mathcal{O}|N\rangle), \tag{8}$$

using the Stokes-like theorem. Let

$$\mathcal{O} = (2-\Delta' f\Delta'^\dagger)f \tag{9}$$

again as in [1], we have for large N

$$\langle N|\mathcal{O}|N\rangle = (\zeta N + x^\mu x^\mu)^{-1} + (\zeta N + x^\mu x^\mu)^{-2}C + O(N^{-3}), \tag{10}$$

where C is some constant $2k \times 2k$ matrix.[3] So

$$\langle N+1|\mathcal{O}|N+1\rangle - \langle N|\mathcal{O}|N\rangle = \frac{-\zeta}{(\zeta N + x^\mu x^\mu)[\zeta(N+1)+x^\mu x^\mu]} + O(N^{-3}). \tag{11}$$

Thus we obtain one part of the topological charge

$$\begin{aligned} Q_1 &= \frac{\zeta}{2\pi}\lim_{N\to\infty}\mathrm{Tr}\int d^2x\,\sum_{n=0}^{N}\langle n|\partial\bar{\partial}[(2-\Delta' f\Delta'^\dagger)f]|n\rangle && (12)\\ &= \frac{-\zeta}{2\pi}\lim_{N\to\infty}\mathrm{Tr}[N\int d^2x\,(\zeta N + x^\mu x^\mu)^{-2} + O(N^{-2})] && (13)\\ &= -k. && (14)\end{aligned}$$

Noting that $\mathcal{O}$ is an ordinary function (matrix) in $x^{3,4}$ and a bounded operator [1] on the Fock space, we immediately work out the other part of the topological charge

$$Q_2 = \frac{\zeta}{8\pi}\mathrm{Tr}\sum_{n=0}^{\infty}\langle n|\int d^2x\,\partial_\mu\partial_\mu[(2-\Delta' f\Delta'^\dagger)f]|n\rangle = 0, \tag{15}$$

---

[3] The non-singularity of each term for arbitrary $x^{3,4}$ in the summation (8) can be analyzed by the same method as in the appendix of [1].

where we have used the Stokes theorem and the following asymptotic as $|x^{3,4}| \to \infty$:

$$f \to \frac{1}{x^\mu x^\mu}. \tag{16}$$

In conclusion, we obtain the topological charge

$$Q = Q_1 + Q_2 = -k, \tag{17}$$

which is expected for an ASD instanton configuration. For an SD instanton on $\mathbf{R}^2_{\mathrm{NC}} \times \mathbf{R}^2$ with instanton number k, we can similarly have

$$Q = k, \tag{18}$$

which completes the answer to the problem of topological charge of noncommutative $U(N)$ instantons.

Acknowledgments

I would like to thank Dr. Jian Dai, Prof. Chuan-Jie Zhu and Prof. Xing-Chang Song for helpful discussions.

References

[1] Y. Tian, C.-J. Zhu and X.-C. Song, *Topological Charge of Noncommutative ADHM Instanton*, Mod. Phys. Lett. A **18** (2003) 1691-1703 [hep-th/0211225].

[2] A. Connes, *Noncommutative Geometry*, Academic Press, New York, 1994.

[3] A. Konechny and A. Schwarz, *Introduction to M(atrix) Theory and Noncommutative Geometry*, Phys. Rept. **360** (2002) 353-465 [hep-th/0012145] [hep-th/0107251].

[4] M. R. Douglas and N. A. Nekrasov, *Noncommutative Field Theory*, Rev. Mod. Phys. 73 (2002) 977, hep-th/0106048.

[5] R. J. Szabo, *Quantum Field Theory on Noncommutative Spaces*, Phys. Rept. **378** (2003) 207-299 [hep-th/0109162].

[6] N. Nekrasov and A. Schwarz, *Instantons on Noncommutative R^4, and (2,0) Superconformal Six Dimensional Theory*, Commun. Math. Phys. 198 (1998) 689, hep-th/9802068.

[7] C.-S. Chu, V. V. Khoze and G. Travaglini, *Notes on Noncommutative Instantons*, Nucl.Phys. B621 (2002) 101, hep-th/0108007.

[8] K.-Y. Kim, B.-H. Lee and H. S. Yang, Phys. Lett. B **523** (2001) 357 [hep-th/0109121].

[9] E. Corrigan, P. Goddard, H. Osborn and S. Templeton, *Zeta Function Regularization And Multi-Instanton Determinants*, Nucl. Phys. B **159** (1979) 469.

[10] Y. Tian and C.-J. Zhu, *Remarks on the noncommutative ADHM construction*, Phys. Rev. D **67** (2003) 045016 [hep-th/0210163].